The Mental Health Guide

Author: US Department of Veteran Affairs VA

Settings: Hospitals

Availability: Free
https://www.patientsafety.va.gov/docs/joe/eps_mental_health_guide.pdf

The Mental Health Guide was developed by a multidisciplinary team comprising of members from the VA National Center for Patient Safety, Nursing, Safety, Environmental Management, and Interior Design to provide guidance and education to the field in relation to determining products suitable for the locked Inpatient Mental Health Environment. The Guide offers recommended products and solutions, is accessed electronically and was designed to be a "living" document updated as

Patient Safety Standards, Materials and Systems Guidelines Recommended by the New York State Office of Mental Health 14th Edition| July 31, 2015 Author: NYS-OMH Settings: Inpatient psychiatric units Availability: Free https://www.omh.ny.gov/omhweb/patient_safety_standards/guide.pdf	The purpose of the New York office of mental health environmental guide is to provide a selection of materials, fixtures, and hardware that the NYS-OMH has reviewed and supports for use within inpatient psychiatric units throughout New York State. While installation of these products will not eliminate all risks, the items selected represent styles and properties of products that help lower patient risk while on an inpatient psychiatric unit. Many of the items in this document represent the current state of the art and it is anticipated that additional or more effective products will continue to be developed. The NYS-OMH intends to periodically update these guidelines to keep current with these changes but notes that hospitals also have an obligation to continue to review products that will assist in this goal. Utilization of any of these products is not mandatory. This document is not intended to provide guidance for outpatient psychiatric facilities. Recommending Organizations: 1. New York State Office of Mental Health Evidence/ Development: Review and evaluation by NY-OMH

Suicide Prevention Resources to support Joint Commission Accredited organizations implementation of NPSG 15.01.01	
EP2 - Validated / Evidence -Based Screening Tools	
Tools	Brief Description

Settings: General, He althcare

Population: All ages

Availability: Free
K W W S V F V V U V F R O X

This triage guide shows how some different types of programs are using the worrisome answers to guide clinical decision making (e.g.. does the patient require 1:1 observation or a psychiatrist to consult?)

The triage model embeds the Columbia Protocol into the Electronic Health Record (EHR) and provides alerts for high risk answers.
7 K H U H L V Q R F R V W R U O L F H Q V H U H T X L U H G I R
providers, to put the Columbia 3 U R W R F R O W R R O V L Q W R (+

Rec ommending Organizations:

1. National Institute of Health NIH
2. Substance Abuse and Mental Health Service Administration SAMHSA
3. National Action Alliance for Suicide Prevention (Action Alliance)
4. Department of Defense
5. CDC National Center for Injury Prevention and Control
6. United States Food and Drug Administration FDA
7. Zero Suicide Initiative http://zerosuicide.sprc.org/

Evidence / Development:
The Columbia Lighthouse Project/Center for Suicide Risk
A29p4(n)2.L33.1(29p4(n(s)-tn2.8(.9 459.4.98 0.3(c)-0.6(.)1 0 2f 0.E>BDC 1 s

(All PHQ, GAD-7 screeners and translations are downloadable from this website and no permission is required to reproduce, translate, display or distrib ute them).	4. Zero Suicide Initiative http://zerosuicide.sprc.org/ Evidence: Gregory E. S et al. Does Response on the PHQ-9 Depression Questionnaire Predict Subsequent Suicide Attempt or Suicide Death? Psychiatric Services 64:1195–1202, 2013; doi: 10.1176/appi.ps.201200587) Rossom RC, Coleman KJ, Ahmedani BK, et al. Suicidal Ideation Reported on the PHQ9 and Risk of Suicidal Behavior across Age Groups. Journal of affective disorders . 2017;215:77-84. doi: 10.1016/j.jad.2017.03.037. Does Suicidal Ideation as measured by the PHQ-9 Predict Suicide Among VA Patients? Samantha A. Louzon, Robert Bossarte, John F. McCarthy, and Ira R. Katz Psychiatric Services 2016 67:5, 517-522
Suicide Behavior Questionnaire- Revised (SBQ -R, Osman et al., 2001)	SuicideB(e)-5.3(h)445(a)243(v)0.7(i)0.8(o)4.1(r)-2.6(Qa)2.3u Revised (SB)-165 RQf i(i)-3.1pCiici2 5165onre

Population: ages 13-18

s-

Suicide risk screening in pediatric hospitals: Clinical pathways to address a global health crisis

Brahmbhatt, Khyati et al. Psychosomatics (2018)

https://www.psychosomaticsjournal.com/article/S0033-3182(18)30429-8/abstract

This paper details the first interdisciplinary and international effort to generate Clinical Pathways (CPs) for pediatric suicide risk screening in general hospital settings.

The Clinical Pathway was created as a guide for hospitals worldwide to improve youth suicide risk screening and implementation of appropriate next steps. The Pathway includes the use of the Ask Suicide-Screening Questions (ASQ) (brief primary screener) and the Columbia Suicide Severity Rating Scale (C-SSRS) or the ASQ Brief Suicide Safety Assessment (secondary screeners) for screening and risk stratification of suicidality in children and adolescents in medical settings (14-17).

The publication includes 4 appendices:

- The introductory document (Appendix A) is intended to help orient providers, managers, and administrators in a variety of disciplines and specialties to the pathway.

- The flow diagrams (Appendix B: 1-3) visually depict the steps in the clinical pathways for suicide risk screening in the ED (Appendix B.1) and in the pediatric inpatient medical/surgical setting (Appendix B.2). Both pathways describe a similar 3-tiered screening process. Further, a brief suicide risk screening for the C-SSRS was created for hospitals that may already be using this scale (Appendix B.3).

- The text document (Appendix C) contains a narrative description of the pathway that is to be used side-by-side with the flow diagrams by individuals or institutions implementing a pediatric

Settings: Primary Care, Behavioral Health, Emergency Departments Availability: Free https://theactionalliance.org /resource/recommended- standard -care	be adopted; and, provides information on resources that are available to make care safer and better.

ED- SAFE Study Materials

The Patient Safety Screener (PSS-

Suicide Prevention Resources to support Joint Commission Accredited organizations implementation of NPSG 15.01.01	
EP3,4 - Validated/ Evidence- Based Suicide Risk Assessment Tools	
Tools	Brief Description
Columbia -Suicide Severity Rating Scale (C- SSRS) Risk Assessment version SAFE- T with C -SSRS Author: Columbia University, the University of Pennsylvania, and the University of Pittsburgh —supported by the National Institute of Mental Health (NIMH) Settings: All Population: all ages and special populations in different settings Availability: Free	The C-SSRSRisk Assessment version can be used as a suicide assessment tool following the use of one of the screening toolsSee http://cssrs.columbia.edu/documents/risk-assessment- page/. The risk assessment version provirsirisk as0 as0 /P <</MCID 48 >>BDC 1 scn 59

to 2. The ratings are then summed to yield a total score, which ranges from 0 to 38. Individual items assess characteristics such as wish to die, desire to make an active or passive suicide attempt, duration and frequency of ideation, sense of control over making an attempt, number of deterrents, and amount of actual preparation for a contemplated attempt. The SSI-W takes approximately 10 minutes to administer. (extract from Brown 2003 , pg 7).

Although the SSI-W has been used less frequently than the SSI, the reliability and validity of this measure have been established.

Evidence:

Brown, G. (2003). A review of suicide assessment measures for intervention research with adults and older adults. Bethesda, MD: National Institute of Mental Health. https://go.edc.org/Brown2003

Beck, A. T., Brown, G. K., & Steer, R. A. (January 01, 1997).

	Evidence: Brown, G. (2003). A review of suicide assessment measures for intervention research with adults and older adults. Bethesda, MD: National Institute of Mental Health. https://go.edc.org/Brown2003 Kreuze E and Lamis D 2017 A Review of Psychometrically Tested Instruments Assessing Suicide Risk in Adults OMEGA—Journal of Death and Dying 2018, Vol. 77(1) 36-90

EP6 - Evidence -based resources for sai142 Tc -baseion142 Tc -ba1r sfi142 uCID z 1.1(r-4(s)-1.8 64 7 7 678 0

F R Q M X Q F W L R Q Z L W K R W K H U L P S R U W D Q W F R P
assessment,appropriate psychopharmacologic treatment,
psychotherapy and hospitalization.

Evidence:
Stanley B, Brown GK, Brenner LA, et al. Comparison of the Safety
Planning Intervention w ith Follow -up vs Usual Care of Suicidal
Patients Treated in the Emergency Department. JAMA
Psychiatry. 2018;75(9):894–900.
doi:10.1001/jamapsychiatry.2018.1776

& X U U L H U * O H Q Q : H W D O Rationale and study protocol fo
intervention: Safety planning and structured follow-up among
veterans at risk for suicide and discharged from the emergency
department Contemporary Clinical Trials, Volume 43, 179 – 184

Barbara Stanley, Gregory K.Brown, Glenn W.

Counseling on Access to Lethal Means (CALM)

Populations: Adults, Youth
Settings: All

Author: Online version - Suicide Prevention Resource Center (SPRC)

Availability: Online - Free

https://training.sprc.org/enrol/index.php?id=3

Counseling on Access to Lethal Means (CALM) is a free, online course from the Suicide Prevention Resource Center about how to reduce access to the methods people use to kill themselves. It covers who needs lethal means counseling and how to work with people at risk of suicide—and their families—to reduce access. While this course is primarily designed for mental health professionals, others who work with people at risk for suicide, such as health care providers and social service professionals, may also benefit. This online course was developed in collaboration with Catherine Barber, director of the Means Matter Campaign at the Harvard Injury Control Research Center, and Elaine Frank, a co-developer of the original in-person CALM workshop.

Recommending Organizations:

1. Suicide Prevention Resource Center
2. SAMHSA
3. Zero Suicide Initiative http://zerosuicide.sprc.org/toolkit/engage#quicktabs-engage=1
4. Harvard Injury Control Research Center, Means Matter

Evidence:

Johnson, R.M., Frank, E.M., Ciocca, M., & Barber, C.W. (2011). Training mental health care providers to reduce at-risk patients' access to lethal means of suicide: Evaluation of CALM Project. Archives of Suicide Research, 15(3), 259-264. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/21827315

Zal

Death by Suicide Within 1 Week of Hospital Discharge: A Retrospective Study of Root Cause Analysis Reports.

[redacted]

Authors: Riblet N, Shiner B, Watts, BV, Mills P, Rusch B, Hempbill RR

[redacted]

Setting: Inpatient mental health unit

Availability: PUBMED

[redacted]

ubmed/28511191

General s uicide reduction tools	
Tool	Brief Description

The Zero Suicide Toolkit

Avai lability: Free